

PRODUCT SPECIFICATIONS

Date: 2015-01-22

PRODUCT NAME	:	Anti-h D-Dimer 1405 SPTN-5
PRODUCT SPECIFICITY	:	Antibody recognizes human D-Dimer
PRODUCT CODE	:	100480
PRODUCT BUFFER	:	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN ₃ as a preservative
SHELF LIFE AND STORAGE	:	Shelf life unspecified, storage at 2-8 °C
ANALYTE DESCRIPTION	:	D-Dimer (DD) is fibrin degradation product created during fibrinolysis when plasmin degrades the fibrin clot. Pulmonary embolism (PE) is challenging diagnosis because most lung scans are inconclusive and pulmonary angiography is an invasive procedure associated with complications. About 70 % of PE are precipitated by deep vein thrombosis (DVT). The DD test is extremely helpful in excluding PE. DD is more sensitive for the diagnosis of proximal DVT than for the diagnosis of distal DVT. DD is also valuable for monitoring patients during and after anticoagulant treatment for recurrent DVT.

PARAMETERS TESTED FROM EACH LOT

PRODUCT APPEARANCE	:	Clear liquid, may precipitate during storage which does not alter the product specifications. Precipitate can be removed by centrifugation or filtering.
PRODUCT CONCENTRATION	:	5.0 mg/ml (+/- 10 %)
PRODUCT ACTIVITY	:	80-120 % compared to reference in an IFMA-test
IEF-RANGE	:	5.8 – 6.7
PURITY	:	≥ 95 %

PARAMETERS DETERMINED ONLY DURING PRODUCT R&D PHASE

CLASS AND SUBCLASS	:	IgG1
ASSOCIATION CONSTANT	:	3.5 x 10 ⁵ 1/Ms
DISSOCIATION CONSTANT	:	does not dissociate
AFFINITY CONSTANT	:	N/A
DETERMINATION METHOD	:	SPR analysis (BioRad ProteOn XPR36)
ANTIGEN	:	FDP-D-Dimer (Chrysal Chem, lot#SDD-1)
CROSS-REACTIVITIES	:	No cross-reactivity against human Fibrinogen, others not tested.

EPITOPE : N/D

EPITOPE GROUP : -

Two antibodies binding to the same, or closely located epitopes, belong to the same group and hence cannot be used as a pair in a sandwich assay. Epitope group numbering does not give any detailed information where the epitope is located.

PAIR RECOMMENDATIONS : **SOLID** **LABEL**

Please note that pair recommendations are based on results obtained in our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations should be taken only as a directive.

PRODUCT STABILITY : **TEMPERATURE, DAYS** **RESULT**

Please note that the shelf life given on page one is based on real time stability testing at +2-8 °C in the product buffer.

-70 °C, 21 days	OK
-20 °C, 21 days	OK
+4 °C, 21 days	OK
+35 °C, 7 days	OK
+35 °C, 21 days	OK, charge changes
+45 °C, 3 days	OK
+45 °C, 7 days	OK, charge changes

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. The maximum duration of the test is 21 days, except for the +45 °C only 7 days.

MISCELLANEOUS : -

REFERENCES : -

Legal disclaimer

MedixMAB monoclonal antibodies meet their specifications if transported, stored and used according to the instructions for use. See also separate *Notes on use and storage of monoclonal antibodies*. If not otherwise agreed in writing all products are sold under *General Delivery Terms*. Other terms may apply if MedixMAB monoclonal antibodies are purchased from an official distributor of Medix Biochemica products. MedixMAB is a registered trademark of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission. Further information on products and methods is available from medixMAB website and extranet.