

PRODUCT SPECIFICATIONS

Date: 2015-05-13

PRODUCT NAME	:	Anti-h Calprotectin 3401 SPTN-5
PRODUCT SPECIFICITY	:	Human Calprotectin S100A9 free subunit
PRODUCT CODE	:	100458
PRODUCT BUFFER	:	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN ₃ as a preservative
SHELF LIFE AND STORAGE	:	Shelf life unspecified, storage at 2-8 °C
ANALYTE DESCRIPTION	:	Calprotectin belongs to the family of S100 proteins and is composed of two subunits S100A8 and S100A9. Increased plasma concentrations are found in response to infections and inflammation. Faecal calprotectin measurement is used for detection of gastrointestinal inflammation or neoplasia and for assessment of inflammatory bowel disease activity and response to treatment.

PARAMETERS TESTED FROM EACH LOT

PRODUCT APPEARANCE	:	Clear liquid, may precipitate during storage which does not alter the product specifications. Precipitate can be removed by centrifugation or filtering.
PRODUCT CONCENTRATION	:	5.0 mg/ml (+/- 10 %)
PRODUCT ACTIVITY	:	80 – 120 % compared to reference in an IFMA-test
IEF-RANGE	:	Does not form focusing bands
PURITY	:	≥ 95 %

PARAMETERS DETERMINED ONLY DURING PRODUCT R&D PHASE

CLASS AND SUBCLASS	:	IgG ₁
ASSOCIATION CONSTANT	:	N/D
DISSOCIATION CONSTANT	:	N/D
AFFINITY CONSTANT	:	N/D
DETERMINATION METHOD	:	N/A
ANTIGEN	:	N/A
CROSS-REACTIVITIES	:	No cross-reactivity with S100A9 / S100A8 complex

EPITOPE : -

EPITOPE GROUP : -

Two antibodies binding to the same, or closely located epitopes, belong to the same group and hence cannot be used as a pair in a sandwich assay. Epitope group numbering does not give any detailed information where the epitope is located.

PAIR RECOMMENDATIONS : **SOLID** **LABEL**

- -

Please note that pair recommendations are based on results obtained in our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations should be taken only as a directive.

PRODUCT STABILITY : **TEMPERATURE, DAYS** **RESULT**

Please note that the shelf life given on page one is based on real time stability testing at +2-8 °C in the product buffer.

-70 °C, 21 days	OK
-20 °C, 21 days	OK
+4 °C, 21 days	OK
+35 °C, 21 days	OK
+45 °C, 7 days	OK

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. The maximum duration of the test is 21 days, except for the +45 °C only 7 days.

MISCELLANEOUS : -

REFERENCES : -

Legal disclaimer

MedixMAB monoclonal antibodies meet their specifications if transported, stored and used according to the instructions for use. See also separate *Notes on use and storage of monoclonal antibodies*. If not otherwise agreed in writing all products are sold under *General Delivery Terms*. Other terms may apply if MedixMAB monoclonal antibodies are purchased from an official distributor of Medix Biochemica products. MedixMAB is a registered trademark of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission. Further information on products and methods is available from medixMAB website.