

PRODUCT SPECIFICATIONS

Date: 2016-03-29

PRODUCT NAME	:	Anti-HRP2 3901 SPTN-5
PRODUCT SPECIFICITY	:	Antibody recognizes <i>Plasmodium falciparum</i> histidine-rich protein 2 (HRP2)
PRODUCT CODE	:	100559
PRODUCT BUFFER	:	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN ₃ as a preservative
SHELF LIFE AND STORAGE	:	Shelf life unspecified, storage at 2-8 °C
ANALYTE DESCRIPTION	:	Malaria is caused by protozoan parasites belonging to the genus <i>Plasmodium</i> . The majority of malaria-related deaths are linked to the species <i>Plasmodium falciparum</i> . Monoclonal antibodies against <i>P. falciparum</i> histidine-rich protein 2 (HRP2) are widely used in malaria rapid diagnostic tests as the parasite secretes substantial amounts of the protein into the host bloodstream.

PARAMETERS TESTED FROM EACH LOT

PRODUCT APPEARANCE	:	Clear liquid, may turn opaque during storage
PRODUCT CONCENTRATION	:	5.0 mg/ml (+/- 10 %)
PRODUCT ACTIVITY	:	80 – 120 % compared to reference in an IFMA-test
IEF-RANGE	:	7.3 – 7.9
PURITY	:	≥ 95 %

PARAMETERS DETERMINED ONLY DURING PRODUCT R&D PHASE

CLASS AND SUBCLASS	:	IgG ₁
ASSOCIATION CONSTANT	:	Not Determined (N/D)
DISSOCIATION CONSTANT	:	N/D
AFFINITY CONSTANT	:	N/D
DETERMINATION METHOD	:	-
ANTIGEN	:	-
CROSS-REACTIVITIES	:	None tested

EPITOPE : N/D

EPITOPE GROUP : -

Two antibodies binding to the same, or closely located epitopes, belong to the same group and hence cannot be used as a pair in a sandwich assay. Epitope group numbering does not give any detailed information where the epitope is located.

PAIR RECOMMENDATIONS : **SOLID** **LABEL**

3902	3901
------	------

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations should be taken only as a directive.

PRODUCT STABILITY : **TEMPERATURE, DAYS** **RESULT**

Please note that the shelf life given on page one is based on real time stability testing at +2-8 °C in the product buffer.

-70 °C, 21 days	OK
-20 °C, 21 days	OK
+4 °C, 21 days	OK
+35 °C, 21 days	OK
+45 °C, 7 days	OK, minor changes in IEF profile

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. The maximum duration of the test is 21 days, except for the +45 °C only 7 days.

MISCELLANEOUS :

REFERENCES :

Legal disclaimer

MedixMAB monoclonal antibodies meet their specifications if transported, stored and used according to the instructions for use. See also separate *Notes on use and storage of monoclonal antibodies*. If not otherwise agreed in writing all products are sold under *General Delivery Terms*. Other terms may apply if MedixMAB monoclonal antibodies are purchased from an official distributor of Medix Biochemica products. MedixMAB is a registered trademark of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission. Further information on products and methods is available from medixMAB website.