

PRODUCT SPECIFICATIONS

Date: 2014-12-19

PRODUCT NAME	:	Anti-h Myoglobin 7005 SPTN-5
PRODUCT SPECIFICITY	:	Antibody recognizes human Myoglobin
PRODUCT CODE	:	100078
PRODUCT BUFFER	:	50 mM Na-citrate, pH 6.5, 0.9 % NaCl, 0.095 % NaN ₃ as a preservative
SHELF LIFE AND STORAGE	:	36 months at 2-8 °C
ANALYTE DESCRIPTION	:	Myoglobin is a single-chain globular protein, containing a heme prosthetic group in the center around which the remaining apoprotein folds. It is the primary oxygen-carrying pigment of muscle tissues. Myoglobin is a sensitive marker for muscle injury.

PARAMETERS TESTED FROM EACH LOT

PRODUCT APPEARANCE	:	Clear liquid, may precipitate during storage which does not alter the product specifications. Precipitate can be removed by centrifugation or filtering.
PRODUCT CONCENTRATION	:	5.0 mg/ml (+/- 10 %)
PRODUCT ACTIVITY	:	80-120 % compared to reference in an IFMA-test
IEF-RANGE	:	5.5 - 6.7
PURITY	:	≥ 95 %

PARAMETERS DETERMINED ONLY DURING PRODUCT R&D PHASE

CLASS AND SUBCLASS	:	IgG ₁
ASSOCIATION CONSTANT	:	N/D
DISSOCIATION CONSTANT	:	N/D
AFFINITY CONSTANT	:	1 x 10 ⁹ l/mol
DETERMINATION METHOD	:	Radioimmunoassay (RIA)
ANTIGEN	:	Myoglobin, Scripps Laboratories (Cat No M0714 Lot 978264)
CROSS-REACTIVITIES	:	No cross-reactivity with Hemoglobin. Others not tested.

EPITOPE : N/D

EPITOPE GROUP : -

Two antibodies binding to the same, or closely located epitopes, belong to the same group and hence cannot be used as a pair in a sandwich assay. Epitope group numbering does not give any detailed information where the epitope is located.

PAIR RECOMMENDATIONS	SOLID	LABEL
	7005	7004

Please note that pair recommendations are based on results obtained in our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations should be taken only as a directive.

PRODUCT STABILITY	TEMPERATURE, DAYS	RESULT
Please note that the shelf life given on page one is based on real time stability testing at +2-8 °C in the product buffer.	-70 °C, 21 days	N/D
	-20 °C, 21 days	OK
	+4 °C, 21 days	OK
	+35 °C, 7 days	OK
	+35 °C, 21 days	OK, but charge changes
	+45 °C, 3 days	OK
	+45 °C, 7 days	OK, but charge changes

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. The maximum duration of the test is 21 days, except for the +45 °C only 7 days.

MISCELLANEOUS : -

REFERENCES : -

Legal disclaimer

MedixMAB monoclonal antibodies meet their specifications if transported, stored and used according to the instructions for use. See also separate *Notes on use and storage of monoclonal antibodies*. If not otherwise agreed in writing all products are sold under *General Delivery Terms*. Other terms may apply if MedixMAB monoclonal antibodies are purchased from an official distributor of Medix Biochemica products. MedixMAB is a registered trademark of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission. Further information on products and methods is available from medixMAB website.